

Confinement from a massive scalar in QCD

Rainer Dick

*Sektion Physik der Ludwig-Maximilians-Universität München
Theresienstr. 37, 80333 München, Germany*

Abstract: A model is introduced with a massive scalar coupling to the Yang-Mills term in four-dimensional gauge theory. It is shown that the resulting potential of colour sources consists of a short range Coulomb interaction and a long range confining part. Far away from the source the scalar vanishes $\sim r^{-1}$ while the potential diverges linearly $\sim r$. Up to an N_c -dependent factor of order 1 the tension parameter in the model is gmf , where m denotes the mass of the scalar and f is a coupling scale entering the scalar-gluon coupling.

1. Recently it was observed that a string inspired coupling of a massless dilaton to gauge fields yields a linearly increasing vector potential from pointlike colour sources, if a logarithmic divergence of the dilaton at infinity is permitted [1, 2]. This motivated me to construct a direct coupling of a massive scalar to chromo-electric and magnetic fields subject to the requirement that the Coulomb problem still admits an analytic solution, but now with a mass term. The model for the scalar-gluon coupling that emerged from this endeavour is

$$\mathcal{L} = -\frac{1}{4} \frac{\phi^2}{f^2 + \beta\phi^2} F_{\mu\nu}^j F^{\mu\nu}_j - \frac{1}{2} \partial_\mu \phi \cdot \partial^\mu \phi - \frac{1}{2} m^2 \phi^2, \quad (1)$$

where $0 \leq \beta \leq 1$ is a parameter and f is a mass scale characterizing the strength of the scalar-gluon coupling.

To analyze the Coulomb problem in this theory we consider a pointlike colour source, which in its rest frame is described by a current

$$j_i^\mu = g\delta(\mathbf{r})C_i\eta^\mu_0.$$

Here $1 \leq i \leq N_c^2 - 1$ is an $\mathfrak{su}(N_c)$ Lie algebra index and $C_i = \zeta^+ \cdot X_i \cdot \zeta$ is the expectation value of the $\mathfrak{su}(N_c)$ generator X_i for a normalized spinor ζ in colour space. These expectation values satisfy

$$\sum_{i=1}^{N_c^2-1} C_i^2 = \frac{N_c - 1}{2N_c},$$

and the equations of motion for the scalar and gluons emerging from this source follow as in [1]:

$$\nabla \cdot \left(\frac{\phi^2}{f^2 + \beta\phi^2} \mathbf{E}_i \right) = gC_i\delta(\mathbf{r}), \quad (2)$$

$$\Delta\phi = m^2\phi - \frac{f^2\phi}{(f^2 + \beta\phi^2)^2} \mathbf{E}^i \cdot \mathbf{E}_i, \quad (3)$$

and $\nabla \times \mathbf{E}_i = 0$ implies existence of chromo-electric potentials $\mathbf{E}_i = -\nabla\Phi_i$.

Clearly, the solution to the Coulomb problem (2,3) includes the case of inertial motion of the colour source through a mere Lorentz boost.

Eq. (2) or more generally its analog for an arbitrary spherically symmetric colour density yields for the fields outside the density

$$\frac{\phi^2}{f^2 + \beta\phi^2} \mathbf{E}_i = \frac{gC_i}{4\pi r^2} \mathbf{e}_r,$$

and inserting this relation in (3) yields

$$\Delta\phi = m^2\phi - \frac{\mu^2}{r^4\phi^3}, \quad (4)$$

where the abbreviation

$$\mu = \frac{gf}{4\pi} \sqrt{\frac{N_c - 1}{2N_c}}$$

was used.

Substituting $y(r) = r\phi(r)$ in (4) and multiplying by dy/dr yields the first integral

$$\left(\frac{dy}{dr}\right)^2 = m^2y^2 + \frac{\mu^2}{y^2} + 2K.$$

This can readily be solved for arbitrary integration constant K , but the boundary condition $\lim_{r \rightarrow \infty} \phi(r) = 0$ uniquely determines $K = -m\mu$. This yields

$$y^2(r) = \frac{\mu}{m} + \left(y_0^2 - \frac{\mu}{m}\right) \exp(-2mr).$$

Therefore, the scalar field emerging from the pointlike colour source is

$$\phi = \pm \frac{1}{r} \sqrt{\frac{\mu}{m} + \left(y_0^2 - \frac{\mu}{m}\right) \exp(-2mr)}, \quad (5)$$

while the chromo-electric potentials consist of a short range Coulomb and a long range confining part:

$$\Phi_i = \beta \frac{gC_i}{4\pi r} - fC_i \sqrt{\frac{N_c}{2(N_c - 1)}} \ln \left(\exp(2mr) - 1 + \frac{m}{\mu} y_0^2 \right). \quad (6)$$

At large distance the scalar field vanishes $\sim r^{-1}$, while the chromo-electric potential yields linear confinement, if applied in the framework of a reduced Salpeter or no-pair equation [3].

2. Eq. (6) implies that an (anti-)quark with colour orientation ζ_q , $\zeta_q^+ \cdot \zeta_q = 1$, in the field of a source of colour ζ_s (a heavy quark) sees a potential

$$V(r) = \pm \left(|\zeta_s^+ \cdot \zeta_q|^2 - \frac{1}{N_c} \right) \left[\beta \frac{g^2}{4\pi r} - gf \sqrt{\frac{N_c}{2(N_c - 1)}} \ln \left(\exp(2mr) - 1 + \frac{m}{\mu} y_0^2 \right) \right], \quad (7)$$

with the upper sign holding for quark-quark interactions and the lower sign applying to quark-anti-quark interactions. The colour factor $|\zeta_s^+ \cdot \zeta_q|^2 - \frac{1}{N_c}$ defines a double cone

around the direction of ζ_s . This double cone has an angle $\tan \theta_c = \sqrt{N_c - 1}$ against the symmetry axis specified by ζ_s , and separates domains of attraction from domains of repulsion: Quarks of colour ζ_q in the double cone are repelled and anti-quarks are attracted, while quarks with colour outside the cone are attracted and anti-quarks are repelled.

Potentials with an $1/r$ singularity at short distances and linear behaviour at large distances have been very successfully applied in the investigation of the quarkonium spectrum, see e.g. [4] for two of the classical references in the field. Phenomenological tension parameters in heavy–light meson systems are of order $\sigma \simeq (430 \text{ MeV})^2$ [5], and in the present model the tension would be determined by the mass and coupling scale of the scalar field according to $\sigma \simeq gmf$.

Eqs. (6,7) indicate that direct couplings of scalar fields to Yang–Mills terms provide an interesting paradigm for the description of confinement in gauge theories. This might be realized in QCD through a fundamental scalar, or eventually through a low energy effective scalar degree of freedom. The possibility of a fundamental scalar cannot be excluded, since the coupling scale f might be large enough to make such a scalar invisible to present day experiments. On the other hand, there exist scalar resonances in the hadronic spectrum whose rôle has not been understood yet.

On the level of a low energy effective scalar degree of freedom, one might speculate that the Lagrangian (1) with $\beta = 0$ is realized in the low energy regime through a QCD dilaton coupling to the trace anomaly, while at high energies the standard QCD Lagrangian would apply. Such a picture could be motivated, if one combines the old idea of scalar meson dominance of the trace of the energy–momentum tensor [6] with the QCD trace anomaly, as in [7]. Since the trace anomaly is proportional to the gluon condensate this could also justify the scalar gluon coupling in (1) with $\beta = 0$, and eq. (7) then tells us how the dilaton changes the quark interaction potential. A disadvantage with this picture concerns the disappearance of the one gluon exchange term in the low energy regime.

In conclusion, the derivation of (7) provides a challenge to monopole condensation as a mechanism for quark confinement, and it seems well justified to dedicate some more efforts to the investigation of phenomenological aspects of scalar–gluon couplings of the kind described in (1).

Acknowledgement: I would like to thank my colleagues in Munich for continuous encouragement and support.

References

- [1] R. Dick, Phys. Lett. B 397 (1997) 193.
- [2] R. Dick, Phys. Lett. B 409 (1997) 321, Fortschr. Phys. 45 (1997) 537.
- [3] J. Sucher, Phys. Rev. A 22 (1980) 348, Phys. Rev. D 51 (1995) 5965;
M.G. Olsson, S. Veseli and K. Williams, Phys. Rev. D 51 (1995) 5079, D 53 (1996) 4006.
- [4] E. Eichten, K. Gottfried, T. Kinoshita, K.D. Lane and T.M. Yan, Phys. Rev. D 21 (1980) 203;
W. Buchmüller and S.-H. H. Tye, Phys. Rev. D 24 (1981) 132.
- [5] S. Godfrey and R. Kokoski, Phys. Rev. D 43 (1991) 1679;
S.N. Gupta and J.M. Johnson, Phys. Rev. D 51 (1995) 168;
D. Scora and N. Isgur, Phys. Rev. D 52 (1995) 2783;
S. Veseli and M.G. Olsson, Phys. Lett. B 393 (1997) 84.
- [6] G. Mack, Nucl. Phys. B 5 (1968) 499;
P.G.O. Freund and Y. Nambu, Phys. Rev. 174 (1968) 1741;
H.A. Kastrup, Nucl. Phys. B 15 (1970) 179;
M. Dal Cin and H.A. Kastrup, Nucl. Phys. B 15 (1970) 189;
J. Ellis, Nucl. Phys. B 22 (1970) 478, B 26 (1971) 536;
J. Wess, in: Scale and Conformal Symmetry in Hadron Physics, ed. R. Gatto (John Wiley & Sons, New York, 1973) pp. 31–41.
- [7] M. Bando, K.-i. Matumoto and K. Yamawaki, Phys. Lett. B 178 (1986) 308;
E. Halyo, Phys. Lett. B 271 (1991) 415.